

## Adult Distal Radius Fracture Management

A. Bobby Chhabra, MD 

Baris Yildirim, MD 

### ABSTRACT

Distal radius fractures (DRFs) are some of the most commonly encountered fractures, and the incidence is increasing. Optimal treatment remains controversial. Critical evaluation of the initial imaging is necessary to recognize fracture characteristics and stability. The fracture pattern, injury mechanism, soft-tissue injury, patient characteristics, and surgeon preference are generally taken into consideration when choosing the most appropriate modality. Volar plating has become the workhorse of surgical DRF management but is not without complications. The surgeon should be comfortable using a wide variety of techniques to customize the fixation to the fracture pattern. Recognition of potential dangers and use of intraoperative imaging techniques can mitigate complications. Goals of rehabilitation after the initial treatment of DRF include regaining motion, strength, and ultimately function while managing pain.

**D**istal radius fractures (DRFs) are some of the most commonly encountered fractures. In the United States, DRF accounts for nearly 17% of all fractures seen in the emergency department with increasing incidence over the past few decades.<sup>1</sup> Optimal treatment is controversial. Here, we present a thorough review of the current literature regarding the management of DRFs.

### Initial Imaging

Management of DRFs is fundamentally guided by radiographs. Radiographic parameters to measure include radial height, radial inclination, volar tilt, ulnar variance, and the teardrop angle; other radiographic factors to assess include intra-articular extension, articular step-off, dorsal comminution, and associated ulnar fracture (Figure 1).<sup>2</sup> The surgeon's recognition of these features guides treatment by gaining a more thorough understanding of the injury pattern, thereby predicting fracture stability.

CT can be used as an additional tool in preoperative evaluation. Because of the cost and radiation associated with its usage, CT scans are typically recommended to evaluate surgical candidates for the extent of joint comminution, gapping, or depression, especially when these factors are not clear with appropriate radiographs (Figure 2). Ideally, CT scans are obtained after initial closed reduction and immobilization to better define the radiographic

From the Department of Orthopaedic Surgery, University of Virginia, Charlottesville, VA.


Correspondence to Dr. Chhabra: ac2h@hscmail.mcc.virginia.edu

Neither of the following authors nor any immediate family member has received anything of value from or has stock or stock options held in a commercial company or institution related directly or indirectly to the subject of this article: Chhabra and Yildirim.

*J Am Acad Orthop Surg* 2021;29:e1105-e1116

DOI: 10.5435/JAAOS-D-20-01335

Copyright 2021 by the American Academy of Orthopaedic Surgeons.

**Figure 1**

PA and lateral radiographs showing measurements of radial height (A), radial inclination (B), ulnar variance (C), volar tilt (D), and teardrop angle (E).

features. Although CT scans can identify articular step-offs and gapping better than plain radiography, their influence in decision-making and patient outcomes is unclear.<sup>3</sup>

## Treatment

A wide variety of nonsurgical and surgical treatment options exist for DRF. Nonsurgical techniques include closed reduction with initial splint or cast immobilization. Surgical techniques include closed reduction with percutaneous pinning; external fixation; dorsal bridge plating; and open reduction with internal fixation (OR-IF) using volar, dorsal, or “fragment-specific” fixation. The fracture pattern, injury mechanism, soft-tissue injury, patient characteristics, and surgeon preference are generally taken into consideration when choosing the most appropriate modality.

One systematic approach is to separate the wrist into the radial, intermediate, and ulnar columns, as described by Rikli and Regazzoni.<sup>4</sup> Rhee et al described a treatment algorithm through the assessment of these individual columns and the pedestal, the metadiaphyseal distal radius, which supports the radial and intermediate columns. For complex DRF, their sequence of wrist

construction is anatomic reduction, fixation of the intermediate column to the pedestal, restoration of the radial column, and any necessary management of ulnar column injuries to stabilize the distal radioulnar joint (DRUJ).<sup>5</sup>

## Closed Management

A common initial practice with DRFs is obtaining reduction through closed manipulation and immobilization. For displaced fractures, recreation of the fracture mechanism with axial traction often leads to acceptable alignment. Two primary methods for closed reduction exist and have been evaluated in the literature: manual traction and finger trap traction. Both methods assist the provider to manipulate and reduce the fracture appropriately by restoring the radial length. Sosborg-Wurtz et al<sup>6</sup> conducted a recent systematic review of the two methods and noted that reduction by manual traction may better correct volar tilt while finger trap traction may better restore radial length, although these results were not found to be clinically significant. Furthermore, finger trap traction may result in a lower incidence of complex regional pain syndrome (CRPS) and carpal tunnel syndrome and cause less pain during reduction. No consensus exists in the choice of analgesia during reduction as well. A recent meta-analysis found that

**Figure 2**

**A**, PA and lateral radiographs showing intra-articular distal radius fractures. **B**, CT scan better revealing intra-articular displacement.

hematoma blocks provide improved postreduction pain relief compared with procedural sedation and analgesia but did not find a difference in pain during reduction or reduction failure.<sup>7</sup> Fernandez<sup>8</sup> investigated the goal radiographic parameters after conducting closed reduction and proposed accepting the following to avoid symptomatic malunion: dorsal tilt  $\leq 10^\circ$ , radial shortening  $\leq 2$  mm, radial inclination  $\geq 15^\circ$ ,  $\leq 2$  mm articular step-off, congruent DRUJ, and absence of carpal malalignment. A close follow-up of these patients is important to ensure that they maintain acceptable

alignment, with patients usually being seen weekly for the first month after their injury.

Radiographic parameters and patient factors need to be considered to determine the fracture's stability, which is defined as its ability to maintain the radiographic position after acceptable closed reduction and immobilization. One of the most frequently discussed studies regarding criteria to evaluate fracture stability was by Lafontaine et al.<sup>9</sup> They retrospectively reviewed a cohort of 167 DRFs and found that the presence of three or more of the following factors was associated with

fracture instability and secondary displacement: dorsal tilt  $>20^\circ$ , dorsal comminution, intra-articular radiocarpal fracture, associated ulnar fracture, and older than 60 years. The significance of these variables continues to be scrutinized in more recent literature. A more recent review emphasized dorsal comminution, female sex, and older than 60 years as increased risk factors of secondary displacement.<sup>10</sup> Another prospective study of 157 patients found that loss of radial height, loss of radial inclination, and older than 52 years were the most important factors in predicting fracture instability.<sup>11</sup> The AAOS adopted appropriate use criteria in 2013 that identified unacceptable reduction criteria as radial shortening greater than 3 mm, dorsal tilt greater than  $10^\circ$ , and articular displacement greater than 2 mm.<sup>12</sup> However, without higher levels of evidence, there will likely be no consensus agreement among surgeons for criteria of treatment.

Insufficient evidence exists for the ideal type of immobilization, duration of immobilization, and initiation of therapy with closed management. Although a sugar tong splint to prevent forearm pronation/supination is the most common method of immobilization, this has been challenged. Splints and casts below the elbow are better tolerated by patients but may sacrifice protection against loss of reduction in unstable fractures. Bong et al<sup>13</sup> found similar performance in reduction maintenance between sugar tong and short arm radial gutter splints in a randomized controlled trial of unstable and stable displaced DRFs. Park et al randomly assigned patients older than 55 years with stable DRFs to short arm and long arm casts. Although they found a notable difference in volar tilt at 3 and 6 months, they found no notable difference in functional outcome scores and found more shoulder pain with long arm casts.<sup>14</sup> Regardless of the type of immobilization, shoulder and finger range-of-motion exercise programs should be initiated immediately after immobilization to minimize stiffness and shoulder pain. Immobilization for 4 to 8 weeks is most common despite insufficient evidence. Early cessation of wrist immobilization risks loss of reduction while attempting to minimize stiffness. In the elderly, stiffness seems to be more common after surgical treatment.<sup>15</sup>

Special consideration is given to the elderly patient with osteoporosis. Although older age has been shown to be a predictor of malunion and instability, patients in this age range seem to have acceptable functional outcomes and satisfaction, regardless of malalignment or loss of reduction radiographically.<sup>16</sup> DeGeorge et al<sup>15</sup> found that nonsurgical management was associated with a notable decrease in 1-year complications for patients in

this cohort. Therefore, this cohort may require separate guidelines for acceptable radiographic alignment and surgical indications. Kodama et al evaluated 52 active and healthy patients specifically of 60 years or older treated with closed management. They found that volar tilt and ulnar variance were the most notable radiographic parameters affecting clinical outcomes in this cohort.<sup>17</sup>

### Closed Reduction and Percutaneous Pinning

Percutaneous pinning can be a useful tool in the fixation of DRFs. Because Kirschner wire fixation is not load-bearing and offers minimal protection against shortening, its use is limited to assisting with initial reduction, unstable DRFs without significant intra-articular involvement, or in conjunction with other modes of fixation that maintain length.

Kapandji<sup>18</sup> first described intrafocal pinning of extra-articular DRFs in 1976, where two Kirschner wires were directed into the fracture site to lever displaced DRFs to improve alignment and maintained to create a buttress effect. Although Kapandji advocated for no additional immobilization postoperatively, this method is typically augmented with additional forms of stability, such as an addition of another wire, or postoperative plaster, or cast immobilization. The ideal duration of postoperative immobilization after percutaneous pinning is unclear.

Percutaneous fixation and ORIF do not seem to have a notable difference in pain or function in the long term. However, when acceptable alignment cannot be achieved through closed or minimal open manipulation, open reduction and internal fixation should be pursued. Although it remains a cost-effective procedure for treating unstable DRF, a recent meta-analysis found that compared with volar plating, closed reduction with percutaneous pinning had a markedly higher incidence of superficial wound (pin tract) infections and CRPS while having lower outcome scores.<sup>19</sup> The technique of burying pins may reduce pin tract infections, although this method may necessitate return to the operating room for wire removal.

### Arthroscopy-assisted Fixation

The use of arthroscopy and its effect on patient outcomes have been debated to supplement the previously mentioned fixation strategies. It can be used to identify suspected associated ligamentous injury, assist with anatomic reduction of intra-articular fractures, and remove intra-articular fracture hematoma and debris. Yamazaki et al<sup>20</sup> found in their randomized controlled trial of 74 patients that arthroscopic assistance did not

improve functional outcomes or radiographic outcomes for unstable intra-articular DRF. Selles et al<sup>21</sup> conducted a randomized controlled trial of 57 patients and found that arthroscopy to remove intra-articular hematoma and debris did not lead to better outcomes with the 1-year follow-up. They also found soft-tissue injuries to the TFCC, SL ligament, and Lister tubercle (LT) ligament injuries in all their patients, but these ligamentous injuries diagnosed on arthroscopy did not require additional treatment in their study. Although arthroscopy remains the benchmark for the diagnosis of SL, LT, and TFCC ligamentous injuries, their treatment after DRF also remains controversial.

### External Fixation

External fixation has long been used in the setting of orthopaedic trauma as a means of both temporary and definitive fixation. The placement of wrist spanning external fixators uses the bare area between the brachioradialis and extensor carpi radialis longus on the forearm and the dorsolateral aspect of the second metacarpal in the hand. Although its use in the treatment of DRF has been mostly supplanted with the introduction of internal fixation techniques, it remains a useful tool for intra-articular DRF with notable comminution and soft-tissue injury, especially in patients who do not require accelerated functional outcome. External fixation can maintain length in the setting of notable comminution through ligamentotaxis but has a small effect on fragment reduction and requires subsequent removal. External fixators can also be used as an adjunct to percutaneous pinning or Kapandji intrafocal pinning for unstable extra-articular DRF.<sup>22</sup> A biomechanical study by Wolfe et al<sup>23</sup> noted that the rigidity of the external fixator can be markedly increased with an augmentation using a single transfixion Kirschner wire.

Complications with external fixation include neuropathies, pin tract infections, pin loosening, nonunion, and malunion. Damage to branches of the superficial sensory radial nerve and lateral antebrachial cutaneous nerves can be avoided with good visualization techniques. The avoidance of pin tract infections requires pin-site care and monitoring, although no pin-site care protocol has been shown to be markedly better than another.<sup>24</sup>

### Dorsal Bridge Plating

Dorsal bridge plating techniques are used in cases of severe comminution, soft-tissue injury, osteoporosis, radiocarpal fracture-dislocation, or in the polytraumatized patient because it provides rigid fixation and distraction across the

wrist with earlier ability for weight-bearing through the extremity (Figure 3).<sup>25</sup> This technique provides the following multiple advantages: it facilitates reduction, prevents radial shortening, permits the soft-tissue envelope to mature, and preserves radiocarpal and radioulnar joint stability. Its popularity has grown especially in preference to external fixation with a similar complication and outcome profile to other treatment methods.<sup>26</sup> Bridge plating can also assist in the salvage of DRF nonunions or malunions. However, widespread adoption of this technique has been stifled by the need for subsequent implant removal surgery and concern for postoperative stiffness. Lauder et al showed a decrease in wrist flexion, wrist extension, and grip strength in dorsal bridge plating of the nondominant extremity, whereas seeing only a loss of wrist flexion in dorsal bridge plating of the dominant extremity.<sup>27</sup> This suggests that rehabilitation and use of the extremity after plate removal may recover some of the lost function.

### Dorsal Fixation

The greatest advantage of dorsal plating is the ability to buttress a dorsally displaced DRF. It may also be useful in dorsal shear fractures and dorsal die-punch fractures. However, compared with other methods of DRF fixation, dorsal plating has fallen out of favor because of its association with subsequent extensor tendon irritation or rupture due to direct contact over the implant, with the extensor pollicis longus (EPL) most commonly affected. This may be mitigated by the removal of implant after fracture healing or use of newer generation plates, which are “low profile” with a thickness roughly half of the original dorsal plates. Using the extensor retinaculum or other biomaterial to cover the dorsal implant has been reported to decrease the occurrence of extensor tendon dysfunction.<sup>28</sup> Should tendon rupture occur, primary repair is usually not possible because of substance atrophy, retraction, or fraying of the tendon ends. The extensor indicis pollicis remains the most commonly used for tendon transfer to the EPL in these situations.

### Volar Fixation

Volar plating has become the most common surgical fixation method preferred by clinicians for its lower complication rate compared with other modalities.<sup>29</sup> Volar locking plates are suitable for unstable extra-articular, basic intra-articular fractures, and osteoporotic individuals, affording early motion and relatively low complication rates (Figure 4). Furthermore, the volar incision can be extended distally to just proximal to the wrist crease, as described by Tannan et al,<sup>30</sup> for a

**Figure 3**



**A**, PA and lateral radiographs showing comminuted distal radius fractures. **B**, Open reduction with internal fixation using dorsal bridge plating and percutaneous pinning.

**Figure 4**



**A**, PA and lateral radiographs showing dorsally displaced distal radius fractures. **B**, Internal fixation using volar plate fixation.

concurrent carpal tunnel release. This method provides the following advantages: no need for incisions over the palm or nerve, no nerve manipulation, reduced retractor-related trauma, and improved exposure to the volar radius. However, the volar approach is not without its limitations.

Volar fixation may not be the best technique for unstable volar rim/shear fractures, fractures with notable articular comminution, or DRFs with DRUJ instability. The improper use of volar plating can lead to iatrogenic tendon injuries, loss of reduction, inadequate fixation, intra-articular screw placement, neurovascular complications, and implant failure.<sup>31-38</sup>

Mehling et al noted in their biomechanical study that the greatest construct stability, highest stiffness under axial compression, and highest load to failure were noted when filling locking screws in all distal plate holes in the distal fragment within 3 mm of the subchondral bone. They recommended at least four screws in the distal fragment with at least two of them in the distal row of the plate.<sup>39</sup> Beck et al<sup>33</sup> found an association with volar plate failure and less than 15 mm of volar cortex available for fixation. This is of specific concern during volar plating for the volar rim or shear fractures because this can lead to excessively distal plate placement leading to iatrogenic tendon injury.

Mechanical and vascular compromise secondary to the surgical repair can lead to tendon irritation, adhesion formation, tenosynovitis, and rupture. For example, flexor tendon irritation can occur with plate placement distal to the “watershed line,” the transverse ridge proximal to the articular surface, and distal to the pronator quadratus. This irritation can ultimately lead to attritional rupture, most commonly of the flexor pollicis longus.<sup>27</sup> In a case series by Drobetz and Kutscha-Lissberg of 50 DRFs treated by the volar locking screw plate system, flexor pollicis longus rupture was noted in 12% of patients at a mean of 10 months postoperatively. They further noted that some of their tendon ruptures could be attributed to improper seating of the locking screw heads, leading to prominence and abrasion of the tendon.<sup>40</sup> Patient education to recognize symptoms and continued follow-ups are important because these complications can occur weeks to months after undergoing ORIF. However, the best way to avoid these complications is with careful placement of volar locking plates with completely flush screw heads proximal to the watershed line. Other strategies to prevent tendon compromise include repair of the pronator quadratus, although no notable advantage has been found in the literature. Using shorter

unicortical screws or smooth pegs,  $\geq 75\%$  of the measured length may also limit complications to extensor tendons.<sup>41</sup> These potential issues should be identified before leaving the operating room by the use of fluoroscopy as outlined in the imaging section. If symptomatic tendinitis occurs, implant removal can resolve symptoms in patients who have gone on to union and prevent tendon rupture.

### Fragment-specific Fixation

Fragment-specific fixation, first described by Medoff and Kopylov in 1998, uses low-profile implants to provide the surgeon with an individualized approach to DRF management. Although more technically demanding, fragment-specific fixation can be a useful tool for a wide range of fracture patterns, including volar rim/shear, ulnar corner, dorsal wall, radial styloid, and free intra-articular fragments (Figure 5). Fragment-specific fixation requires identification of these DRF implants, followed by independent fixation with semi-elastic fixation mechanisms to restore articular anatomy without the need for effective thread purchase in small articular fragments.<sup>42</sup> This creates a load-sharing construct with a theoretical advantage to permit earlier range of motion. A biomechanical study by Taylor et al<sup>43</sup> revealed no notable difference in the stiffness of the constructs between fragment-specific fixation and fixed-angle volar plating, implying that they would do similarly with an early range-of-motion protocol. When compared with volar plating for outcomes, Sammer et al found fragment-specific fixation to be less stable and have worse outcomes at 6 months with similar outcomes at 12 months. They found higher complication and revision surgery rates with fragment-specific fixation, although their study was limited by selection bias.<sup>44</sup> A more recent randomized study by Landgren et al<sup>45</sup> of 50 DRFs revealed no statistical difference in patient-reported outcomes between volar plating and fragment-specific fixation at 12 months but found a markedly higher complication rate in the fragment-specific fixation group. Careful patient selection with fragment-specific fixation is important because this technique often requires multiple incisions and plates, which may cause issues with tendon irritation.

### Intraoperative Imaging

In the surgical setting, the ability to recognize potential problems and ensuring adequate reduction using imaging is paramount. PA and lateral views will provide the




**Figure 5**

**A**, PA and lateral radiographs showing a multifragmentary distal radius fracture about the radial, intermediate, and ulnar columns. **B**, Fragment-specific fixation to capture and stabilize the individual pieces.

standard overview of the alignment, length, and volar tilt while exposing the anatomic landmarks for ideal plate placement. Oblique radiographic views can be helpful in

the detection of implant complications. The anatomic wrist PA view, obtained with 11° of volar tilt with the wrist neutral, and the anatomic tilt lateral view,

**Figure 6**




**A,** Intraoperative fluoro images of the anatomic PA and anatomic lateral views showing no intra-articular screw penetration.

obtained with 15° to 30° radial tilt with the wrist in a neutral position, allow the surgeon to visualize potential intra-articular screw penetration in both dorsal and volar plating techniques (Figure 6). In combination, both of these views provide an improved assessment of intra-articular screw penetration compared with the standard PA and lateral views. A 45° pronated oblique

image can be used to visualize both the lunate and scaphoid facets. The dorsal tangential view identifies dorsal screw penetration in volar plating techniques, which can affect extensor tendons. The view is obtained with the wrist in near full flexion and using fluoroscopy to visualize the full height of LT, thereby identifying the greatest distance between LT and EPL groove (Figure 7).

**Figure 7**



**A,** Intraoperative fluoro image showing no dorsal screw penetration.

If implant complications cannot be excluded with standard and oblique imaging, consider “live” fluoroscopic imaging or arthroscopic assistance.

Goals of rehabilitation after the initial treatment of DRF include regaining motion, strength, and ultimately function while managing pain. Postoperative rehabilitation of DRFs may be divided into the following three stages: initial immobilization for pain control and early wound healing, mobilization, and strengthening.<sup>46</sup>

Understanding of the optimal length of immobilization after initial DRF management continues to evolve. Shorter postoperative immobilization times have been found to improve range of motion in the early postoperative period without leading to increased complications.<sup>47,48</sup> Furthermore, an accelerated rehabilitation protocol allowing “immediate” gentle range of motion may be safe and allow for early return to function.<sup>48</sup> However, there is a paucity of literature demonstrating durable benefits to early mobilization.

Whether a home exercise program or supervised therapy after surgical fixation of DRF is more effective is also debated. A prospective randomized study found that formal occupational therapy was comparable with or slightly inferior to physician-directed independent home exercise programs for range of motion and grip strength after volar plating of DRF.<sup>49</sup> Valdes et al found no difference between those in a home exercise program or clinic-based therapy in patients who underwent uncomplicated volar plating. However, in patients with postoperative complications (such as finger stiffness, CRPS, and carpal tunnel syndrome) or with comorbidities (such as osteoarthritis or advanced age), clinic-based therapy may be advantageous.<sup>50</sup>

## Summary

Successful outcomes of treating DRF are multifaceted. It is necessary to achieve appropriate reduction and restoration of native anatomy for the goal of return to mobility, function, work, and/or sport within a reasonable timeframe. Understanding all available modalities and their potential complications will lead to improved success in treatment of each patient’s fracture.

## Acknowledgment

The authors thank David M. Noble, MD for assistance with manuscript preparation.

## References

References printed in **bold type** are those published within the past 5 years.

1. Nellans KW, Kowalski E, Chung KC: The epidemiology of distal radius fractures. *Hand Clin* 2012;28:113-125.
2. Medoff RJ: Essential radiographic evaluation for distal radius fractures. *Hand Clin* 2005;21:279-288.
3. das Gracias Nascimento V, da Costa AC, Falcochio DF, Lanzarin LD, Checchia SL, Chakkour I: Computed tomography’s influence on the classifications and treatment of the distal radius fractures. *Hand (N Y)* 2015;10:663-669.
4. Rikli DA, Regazzoni P: Fractures of the distal end of the radius treated by internal fixation and early function. A preliminary report of 20 cases. *J Bone Joint Surg Br Vol* 1996;78:588-592.
5. **Rhee PC, Medoff RJ, Shin AY: Complex distal radius fractures: An anatomic algorithm for surgical management. *J Am Acad Orthop Surg* 2017;25:77-88.**
6. **Sosborg-Wurtz H, Corap Gellert S, Ladeby Erichsen J, Viberg B: Closed reduction of distal radius fractures: A systematic review and meta-analysis. *EFORT Open Rev* 2018;3:114-120.**
7. **Tseng PT, Leu TH, Chen YW, Chen YP: Hematoma block or procedural sedation and analgesia, which is the most effective method of anesthesia in reduction of displaced distal radius fracture? *J Orthop Surg Res* 2018;13:1-7.**
8. Fernandez DL: Should anatomic reduction be pursued in distal radial fractures? *J Hand Surg* 2000;25:523-527.
9. Lafontaine M, Hardy D, Delince P: Stability assessment of distal radius fractures. *Injury* 1989;20:208-210.
10. Walenkamp MM, Aydin S, Mulders MA, Goslings JC, Schep NW: Predictors of unstable distal radius fractures: A systematic review and meta-analysis. *J Hand Surg Eur Vol* 2016;41:501-515.
11. Tahirian MA, Javdan M, Nouraei MH, Dehghani M: Evaluation of instability factors in distal radius fractures. *J Res Med Sci* 2013;18:892.
12. Hammert WC, Kramer RC, Graham B, Keith MW: AAOS appropriate use criteria: Treatment of distal radius fractures. *J Orthop Surg Res* 2013; 21:506-509.
13. Bong MR, Egol KA, Leibman M, Koval KJ: A comparison of immediate postreduction splinting constructs for controlling initial displacement of fractures of the distal radius: A prospective randomized study of long-arm versus short-arm splinting. *J Hand Surg* 2006;31:766-770.
14. **Park MJ, Kim JP, Lee HI, Lim TK, Jung HS, Lee JS: Is a short arm cast appropriate for stable distal radius fractures in patients older than 55 years? A randomized prospective multicentre study. *J Hand Surg Eur Vol* 2017;42:487-492.**
15. **DeGeorge BR, Jr, Van Houten HK, Mwangi R, Sangaralingham LR, Larson AN, Kakar S: Outcomes and complications in the management of distal radial fractures in the elderly. *J Bone Joint Surg Am* 2020;102: 37-44.**
16. Grewal R, MacDermid JC: The risk of adverse outcomes in extra-articular distal radius fractures is increased with malalignment in patients of all ages but mitigated in older patients. *J Hand Surg Am* 2007;32: 962-970.
17. Kodama N, Takemura Y, Ueba H, Imai S, Matsusue Y: Acceptable parameters for alignment of distal radius fracture with conservative treatment in elderly patients. *J Orthop Sci* 2014;19:292-297.
18. Kapandji A: Internal fixation by double intrafocal plate. Functional treatment of non articular fractures of the lower end of the radius (author’s transl) [in French]. *Ann Chir* 1976;30:903-908.

19. Peng F, Liu YX, Wan ZY: Percutaneous pinning versus volar locking plate internal fixation for unstable distal radius fractures: A meta-analysis. *J Hand Surg Eur Vol* 2018;43:158-167.
20. Yamazaki H, Uchiyama S, Komatsu M, et al: Arthroscopic assistance does not improve the functional or radiographic outcome of unstable intra-articular distal radial fractures treated with a volar locking plate: A randomised controlled trial. *Bone Joint J* 2015;97:957-962.
21. Selles CA, Mulders MA, Colaris JW, van Heijl M, Cleffken BI, Schep NW: Arthroscopic debridement does not enhance surgical treatment of intra-articular distal radius fractures: A randomized controlled trial. *J Hand Surg Eur Vol* 2020;45:327-332.
22. Weil WM, Trumble TE: Treatment of distal radius fractures with intrafocal (kapandji) pinning and supplemental skeletal stabilization. *Hand Clin* 2005;21:317-328.
23. Wolfe SW, Swigart CR, Grauer J, Slade JF III, Panjabi MM: Augmented external fixation of distal radius fractures: A biomechanical analysis. *J Hand Surg Am* 1998;23:127-134.
24. Ktistakis I, Guerado E, Giannoudis PV: Pin-site care: Can we reduce the incidence of infections? *Injury* 2015;46(suppl 3):S35-S39.
25. Hyatt BT, Hanel DP, Saucedo JM: Bridge plating for distal radius fractures in low-demand patients with assist devices. *J Hand Surg Am* 2019;44:507-513.
26. Lauder A, Agnew S, Bakri K, Allan CH, Hanel DP, Huang JI: Functional outcomes following bridge plate fixation for distal radius fractures. *J Hand Surg Am* 2015;40:1554-1562.
27. Tanaka Y, Aoki M, Izumi T, Fujimiya M, Yamashita T, Imai T: Effect of distal radius volar plate position on contact pressure between the flexor pollicis longus tendon and the distal plate edge. *J Hand Surg Am* 2011;36:1790-1797.
28. Rhee PC, Dennison DG, Kakar S: Avoiding and treating perioperative complications of distal radius fractures. *Hand Clin* 2012;28:185-198.
29. Wichlas F, Haas NP, Disch A, Machó D, Tsitsilonis S: Complication rates and reduction potential of palmar versus dorsal locking plate osteosynthesis for the treatment of distal radius fractures. *J Orthop Traumatol* 2014;15:259-264.
30. Tannan SC, Pappou IP, Gwathmey FW, Freilich AM, Chhabra AB: The extended flexor carpi radialis approach for concurrent carpal tunnel release and volar plate osteosynthesis for distal radius fracture. *J Hand Surg Am* 2015;40:2026-2031.e1.
31. Koo SC, Ho ST: Delayed rupture of flexor pollicis longus tendon after volar plating of the distal radius. *Hand Surg* 2006;11:67-70.
32. Failla JM, Koniuch MP, Moed BR: Extensor pollicis longus rupture at the tip of a prominent fixation screw: Report of three cases. *J Hand Surg Am* 1993;18:648-651.
33. Beck JD, Harness NG, Spencer HT: Volar plate fixation failure for volar shearing distal radius fractures with small lunate facet fragments. *J Hand Surg Am* 2014;39:670-678.
34. Rozental TD, Blazar PE: Functional outcome and complications after volar plating for dorsally displaced, unstable fractures of the distal radius. *J Hand Surg Am* 2006;31:359-365.
35. Arora R, Lutz M, Hennerbichler A, Krappinger D, Espen D, Gabl M: Complications following internal fixation of unstable distal radius fracture with a palmar locking-plate. *J Orthop Trauma* 2007;21:316-322.
36. Biemek T, Kusz D, Cielinski L: Peripheral nerve compression neuropathy after fractures of the distal radius. *J Hand Surg* 2006;31:256-260.
37. Dao KD, Venn-Watson E, Shin AY: Radial artery pseudoaneurysm complication from use of AO/ASIF volar distal radius plate: A case report. *J Hand Surg Am* 2001;26:448-453.
38. De Baere T, Lecouvet F, Barbier O: Breakage of a volar locking plate after delayed union of a distal radius fracture. *Acta Orthop Belg* 2007;73:785-790.
39. Mehling I, Müller LP, Delinsky K, Mehler D, Burkhart KJ, Rommens PM: Number and locations of screw fixation for volar fixed-angle plating of distal radius fractures: Biomechanical study. *J Hand Surg Am* 2010;35:885-891.
40. Drobetz H, Kutscha-Lissberg E: Osteosynthesis of distal radial fractures with a volar locking screw plate system. *Int Orthop* 2003;27:1-6.
41. Wall LB, Brodt MD, Silva MJ, Boyer MI, Calfee RP: The effects of screw length on stability of simulated osteoporotic distal radius fractures fixed with volar locking plates. *J Hand Surg Am* 2012;37:446-453.
42. Medoff RJ: Fragment-specific fixation of distal radius fractures. *Oper Tech Orthop Trauma Surg* 2015:75-93.
43. Taylor KF, Parks BG, Segalman KA: Biomechanical stability of a fixed-angle volar plate versus fragment-specific fixation system: Cyclic testing in a C2-type distal radius cadaver fracture model. *J Hand Surg Am* 2006;31:373-381.
44. Sammer DM, Fuller DS, Kim HM, Chung KC: A comparative study of fragment-specific versus volar plate fixation of distal radius fractures. *Plast Reconstr Surg* 2008;122:1441-1450.
45. Landgren M, Abramo A, Geijer M, Kopylov P, Tägil M: Fragment-specific fixation versus volar locking plates in primarily nonreducible or secondarily redisplaced distal radius fractures: A randomized controlled study. *J Hand Surg Am* 2017;42:156-165.e1.
46. Ikpeze TC, Smith HC, Lee DJ, Elfar JC: Distal radius fracture outcomes and rehabilitation. *Geriatr Orthop Surg Rehabil* 2016;7:202-205.
47. Watson N, Haines T, Tran P, Keating JL: A comparison of the effect of one, three, or six weeks of immobilization on function and pain after open reduction and internal fixation of distal radial fractures in adults: A randomized controlled trial. *J Bone Joint Surg Am* 2018;100:1118-1125.
48. Quadlbauer S, Pezzeri C, Jurkowitsch J, et al: Early rehabilitation of distal radius fractures stabilized by volar locking plate: A prospective randomized pilot study. *J Wrist Surg* 2017;6:102-112.
49. Souer JS, Buijze G, Ring D: A prospective randomized controlled trial comparing occupational therapy with independent exercises after volar plate fixation of a fracture of the distal part of the radius. *JBSJ* 2011;93:1761-1766.
50. Valdes K, Naughton N, Burke CJ: Therapist-supervised hand therapy versus home therapy with therapist instruction following distal radius fracture. *J Hand Surg Am* 2015;40:1110-1116.e1.